

Table 2. Confidence intervals (95%) for the amount of moisture lost to evapotranspiration before 43 Kentucky bluegrass entries from data set 18APP2 reach 75, 50, and 25% green cover. Entries are ranked based on their ET₅₀ values.

Rank	Selection	ET ₇₅	ET ₅₀	ET ₂₅
1.	Mystere H2O	6.941 to 7.412	7.830 to 8.172	8.576 to 9.077
2.	A08-2	6.736 to 7.147	7.571 to 7.872	8.281 to 8.721
3.	NA1-17-92	6.515 to 6.814	7.285 to 7.485	7.954 to 8.258
4.	A04-383	6.317 to 6.848	7.177 to 7.537	7.869 to 8.398
5.	PPG-KB1316	6.295 to 6.761	7.123 to 7.431	7.802 to 8.251
6.	Mallard	6.001 to 6.666	6.950 to 7.405	7.703 to 8.340
7.	PPG-KB1304	5.976 to 6.524	6.938 to 7.318	7.743 to 8.271
8.	Blue Devil	5.553 to 6.296	6.738 to 7.281	7.711 to 8.479
9.	Diva	5.943 to 6.379	6.800 to 7.098	7.539 to 7.934
10.	AKB3208	5.728 to 6.275	6.743 to 7.135	7.609 to 8.146
11.	NAI-11-644	5.398 to 6.222	6.619 to 7.229	7.607 to 8.470
12.	Rubix	5.612 to 6.225	6.531 to 6.975	7.297 to 7.880
13.	NAI-17-88	5.073 to 5.842	6.446 to 7.036	7.600 to 8.452
14.	AKB1925	5.224 to 5.796	6.474 to 6.909	7.565 to 8.181
15.	Rockstar	5.232 to 6.100	6.296 to 6.948	7.138 to 8.024
16.	AKB3241(Engeti)	5.521 to 6.017	6.435 to 6.805	7.226 to 7.716
17.	AKB3169	4.354 to 5.357	6.046 to 6.855	7.436 to 8.654
18.	NAI-13-9	5.203 to 5.735	6.223 to 6.633	7.106 to 7.668
19.	NAI-15-84	5.031 to 5.755	6.118 to 6.678	7.019 to 7.794
20.	Mazama	5.351 to 5.712	6.236 to 6.518	7.031 to 7.413
21.	NAI-13-14	4.973 to 5.489	6.117 to 6.520	7.124 to 7.689
22.	KH3492(Tejas)	5.195 to 5.747	6.071 to 6.509	6.808 to 7.409
23.	AKB1892	4.819 to 5.565	5.990 to 6.574	6.963 to 7.783
24.	Rubicon	5.331 to 5.714	6.111 to 6.423	6.797 to 7.226
25.	Midnight	4.906 to 5.505	5.984 to 6.455	6.905 to 7.563
26.	A16-8	4.712 to 5.398	5.800 to 6.345	6.705 to 7.476
27.	NAI-13-132	4.820 to 5.279	5.861 to 6.225	6.778 to 7.295
28.	Zip	4.551 to 5.335	5.696 to 6.321	6.630 to 7.520
29.	NAI-13-1	3.865 to 5.029	5.517 to 6.467	6.831 to 8.249
30.	A15-18	4.818 to 5.300	5.768 to 6.150	6.584 to 7.133
31.	Shamrock	4.594 to 5.250	5.639 to 6.162	6.504 to 7.255
32.	15-105	4.489 to 5.033	5.652 to 6.090	6.666 to 7.296
33.	AKB1162	4.449 to 5.052	5.596 to 6.082	6.579 to 7.278
34.	AKB3348	4.532 to 5.121	5.549 to 6.019	6.402 to 7.081
35.	Ginney II	4.204 to 4.962	5.410 to 6.026	6.407 to 7.299
36.	A15-21	4.198 to 4.916	5.355 to 5.938	6.312 to 7.158
37.	NAI-12-41	3.993 to 4.751	5.222 to 5.843	6.238 to 7.146
38.	Quartz	4.141 to 4.827	5.252 to 5.809	6.171 to 6.982
39.	NAI-14-128	4.037 to 4.663	5.089 to 5.598	5.966 to 6.708

40.	NAI-14-122	3.817 to 4.287	4.866 to 5.254	5.781 to 6.355
41.	NAI-14-187	3.708 to 4.269	4.797 to 5.262	5.725 to 6.415
42.	NAI-14-133	3.416 to 3.911	4.577 to 4.988	5.590 to 6.213
43.	Snap	3.121 to 3.821	4.342 to 4.918	5.346 to 6.232




Figure 2. Confidence intervals (95%) for the moisture lost to evapotranspiration before the 43 Kentucky bluegrass entries from data set 18APP2 reach 75, 50, and 25% green cover.