

NATIONAL TURFGRASS EVALUATION PROGRAM

The National Turfgrass Evaluation Program (NTEP) is designed to develop and coordinate uniform evaluation trials of turfgrass varieties and promising selections in the United States and Canada. Test results can be used by national companies and plant breeders to determine the broad picture of the adaptation of a cultivar. Results can also be used to determine if a cultivar is well adapted to a local area or level of turf maintenance.

Briefly, the NTEP is a self-supporting, non-profit program, sponsored by the Beltsville Agricultural Research Center and the National Turfgrass Federation, Inc. Program policy is made by a policy committee consisting of one member from each of the four (4) Regional Turfgrass Research Committees in the United States, one member from the Lawn Seed Division of the American Seed Trade Association, one member from the United States Golf Association (USGA) Green Section, one member from the Golf Course Superintendents Assoc. of America (GCSAA), one member from the Turfgrass Producers International (TPI), one member from the Turfgrass Breeders Association and an executive director. The program does not make variety recommendations. However, the data from tests can be used by extension specialists and others for making recommendations.

The policy committee is responsible for determining program policy including, (1) requirements for submission of entries, (2) scheduling tests, (3) evaluation methods, (4) selecting standard or control test entries, (5) setting entry fees, (6) coordinating tests in their respective regions, (7) establishing guidelines for publication and data distribution and (8) scheduling committee meetings.

Executive Director - Kevin N. Morris, National Turfgrass Evaluation Program, Inc.

CURRENT POLICY COMMITTEE MEMBERS:

Mr. Bill Dunn, Seed Research of Oregon, Inc.
Mr. Ryan Jeffries, Columbia Seeds, Inc.
Dr. Mike Kenna, USGA Green Section
Dr. Charles Fontanier, Oklahoma State University
Dr. David Kopec, University of Arizona
Mr. Gary Wilbur, Oakwood Sod Farms
Mr. Mark Johnson, Golf Course Superintendents Assoc. of America
Dr. Aaron Patton, Purdue University
Mr. Steve Reid, DLF Pickseed USA
Mr. Steve Rackliffe, University of Connecticut

FOR ADDITIONAL REPORTS OR INFORMATION CONTACT:

Kevin Morris, Executive Director
National Turfgrass Evaluation Program
Beltsville Agricultural Research Center-West
Building 005, Room 307
Beltsville, Maryland 20705
kmorris@ntep.org
www.ntep.org

CONTENTS

2014 National Bentgrass (Fairway/Tee) Test - 2017 data

LOCATIONS SUBMITTING DATA FOR 2017.....	1
NATIONAL BENTGRASS (FAIRWAY/TEE) TEST, 2014	
Entries and Sponsors.....	2
Table A - 2017 Locations, Site Descriptions and Management Practices in the 2014 National Bentgrass (Fairway/Tee) Test.....	3
Table B - Locations and Data Collected in 2017.....	4
Table 1A - Mean Turfgrass Quality Ratings of Bentgrass Cultivars Grown on a Fairway or Tee Grown in Location Performance Index (LPI) Group 1.....	6
Table 1B - Mean Turfgrass Quality Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee Grown in Location Performance Index (LPI) Group 1.....	7
Table 1C - Mean Turfgrass Quality Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee Grown in Location Performance Index (LPI) Group 1.....	7
Table 2A - Mean Turfgrass Quality Ratings of Bentgrass Cultivars Grown on a Fairway or Tee Grown in Location Performance Index (LPI) Group 2.....	8
Table 2B - Mean Turfgrass Quality Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee Grown in Location Performance Index (LPI) Group 2.....	9
Table 2C - Mean Turfgrass Quality Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee Grown in Location Performance Index (LPI) Group 2.....	9
Table 3A - Mean Turfgrass Quality and Other Ratings of Bentgrass Cultivars Grown on a Fairway or Tee Under Traffic Stress at Amherst, MA.....	10
Table 3B - Mean Turfgrass Quality and Other Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee under Traffic Stress at Amherst, MA.....	11
Table 3C - Mean Turfgrass Quality and Other Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee under Traffic Stress at Amherst, MA.....	11
Table 4A - Mean Turfgrass Quality and Other Ratings of Bentgrass Cultivars Grown on a Fairway or Tee under Traffic Stress at Blacksburg, VA.....	12
Table 4B - Mean Turfgrass Quality and Other Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee under Traffic Stress at Blacksburg, VA.....	13

CONTENTS (Continued)

Table 4C - Mean Turfgrass Quality and Other Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee under Traffic Stress at Blacksburg, VA.....	13
Table 5A - Mean Turfgrass Quality and Other Ratings of Bentgrass Cultivars Grown on a Fairway or Tee under Reduced Irrigation at Riverside, CA.....	14
Table 5B - Mean Turfgrass Quality and Other Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee under Reduced Irrigation at Riverside, CA.....	15
Table 5C - Mean Turfgrass Quality and Other Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee under Reduced Irrigation at Riverside, CA.....	15
Table 6A - Mean Turfgrass Quality and Other Ratings of Bentgrass Cultivars Grown on a Fairway or Tee under Reduced Irrigation at West Lafayette, IN.....	16
Table 6B - Mean Turfgrass Quality and Other Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee under Reduced Irrigation at West Lafayette, IN.....	17
Table 6C - Mean Turfgrass Quality and Other Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee under Reduced Irrigation at West Lafayette, IN.....	17
Table 7A - Mean Turfgrass Quality and Other Ratings of Bentgrass Cultivars Grown on a Fairway or Tee under Reduced Irrigation at Logan, UT.....	18
Table 7B - Mean Turfgrass Quality and Other Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee under Reduced Irrigation at Logan, UT.....	19
Table 7C - Mean Turfgrass Quality and Other Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee under Reduced Irrigation at Logan, UT.....	19
Table 8A - Mean Turfgrass Quality and Other Ratings of Bentgrass Cultivars Grown on a Fairway or Tee under Low Temperature/Winter Tolerance at Fargo, ND.....	20
Table 8B - Mean Turfgrass Quality and Other Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee under Low Temperature/Winter Tolerance at Fargo, ND.....	21
Table 8C - Mean Turfgrass Quality and Other Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee under Low Temperature/Winter Tolerance at Fargo, ND.....	21
Table 9A - Genetic Color Ratings of Bentgrass Cultivars Grown on a Fairway or Tee.....	22
Table 9B - Genetic Color Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee.....	23

CONTENTS (Continued)

Table 9C - Genetic Color Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee.....	23
Table 10A- Spring Greenup Ratings of Bentgrass Cultivars Grown on a Fairway or Tee.....	24
Table 10B- Spring Greenup Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee.....	25
Table 10C- Spring Greenup Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee.....	25
Table 11A- Leaf Texture Ratings of Bentgrass Cultivars Grown on a Fairway or Tee.....	26
Table 11B- Leaf Texture Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee.....	27
Table 11C- Leaf Texture Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee.....	27
Table 12A- Fall Density Ratings of Bentgrass Cultivars Grown on a Fairway or Tee.....	28
Table 12B- Fall Density Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee.....	29
Table 12C- Fall Density Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee.....	29
Table 13A- Dollar Spot Ratings of Bentgrass Cultivars Grown on a Fairway or Tee.....	30
Table 13B- Dollar Spot Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee.....	31
Table 13C- Dollar Spot Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee.....	31
Table 14A- Brown Patch (Warm Temperature) Ratings of Bentgrass Cultivars Grown on a Fairway or Tee.....	32
Table 14B- Brown Patch (Warm Temperature) Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee.....	33
Table 14C- Brown Patch (Warm Temperature) Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee.....	33
Table 15A- Anthracnose Ratings of Bentgrass Cultivars Grown on a Fairway or Tee.....	34
Table 15B- Anthracnose Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee.....	35
Table 15C- Anthracnose Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee.....	35

CONTENTS (Continued)

Table 16A- Fall Color (October) Ratings of Bentgrass Cultivars Grown on a Fairway or Tee.....	36
Table 16B- Fall Color (October) Ratings of Creeping Bentgrass Cultivars Grown on a Fairway or Tee.....	37
Table 16C- Fall Color (October) Ratings of Colonial Bentgrass Cultivars Grown on a Fairway or Tee.....	37
Appendix Table- Summary of Turfgrass Quality Ratings of Bentgrass Cultivars Grown on a Fairway or Tee.....	38

A Guide to NTEP Turfgrass Ratings

Introduction

The quality and scientific merit of NTEP data is extremely important. However, the evaluation of turfgrass species and cultivars is a difficult and complex issue. Furthermore, turfgrass evaluation is generally a subjective process based on visual estimates of factors, like genetic color, stand density, leaf texture, uniformity and quality. These factors can not be measured in the same way as other agricultural crops. Turfgrass quality is not a measure of yield or nutritive value. Turfgrass quality is a measure of aesthetics (i.e. density, uniformity, texture, smoothness, growth habit and color), and functional use. The most common way of assessing turfgrass quality is a visual rating system that is based on the turfgrass evaluator's judgement.

General Considerations

Most visual ratings collected on NTEP trials are based on a 1 to 9 rating scale. One is the poorest or lowest and 9 is the best or highest rating. However, a few characteristics, such as winter kill or percent living ground cover, are rated on a percentage basis, again by using the evaluator's judgement. Most disease ratings found in NTEP reports will use the 1-9 scale, 9=no disease except where the evaluator made a judgement of the percentage of disease in each plot. Percent disease data will be found in separate tables and will normally not be included with disease data using the 1-9 scale.

Turfgrass Quality

Turfgrass Quality is based on 9 being outstanding or ideal turf and 1 being poorest or dead. A rating of 6 or above is generally considered acceptable. A quality rating value of 9 is reserved for a perfect or ideal grass, but it also can reflect an absolutely outstanding treatment plot. The NTEP requires quality ratings on a monthly basis. Quality ratings take into account the aesthetic and functional aspects of the turf. Quality ratings are not based on color alone, but on a combination of color, density, uniformity, texture, and disease or environmental stress.

Turfgrass quality ratings are grouped and presented by region, management level, a particular stress (shade, traffic, etc.) and in some cases, by individual location (starting with 2001 data, data from each location will be posted separately as well on the NTEP web site, <http://www.ntep.org>). Also available now is a summary table (Appendix) in the back of this report. This summary table includes various statistical measures not previously compiled for NTEP reports. For an explanation of this table and these changes, please go to the NTEP web site at <http://www.ntep.org/pdf/grandmean.mem.pdf>.

Other Ratings

More detailed information on the ratings of specific characteristics can be found on the NTEP web site at <http://www.ntep.org/reports/ratings.htm>.

2014 NATIONAL BENTGRASS TEST
(Fairway/Tee)

LOCATIONS SUBMITTING DATA FOR 2017

State	Location	Code
California	Riverside (Reduced Irrigation)	CA3
Illinois	Urbana	IL1
Indiana	West Lafayette (Reduced Irrigation)	IN1
Iowa	Ames	IA1
Kansas	Manhattan	KS1
Kentucky	Lexington	KY1
Massachusetts	Amherst (Traffic)	MA1
Maryland	College Park	MD1
Michigan	East Lansing	MI1
Missouri	Columbia	MO1
New Jersey	North Brunswick	NJ1
North Dakota	Fargo (Winter Tolerance)	ND1
Utah	Logan (Reduced Irrigation)	UT1
Virginia	Blacksburg (Traffic)	VA2
Washington	Pullman	WA1

2014 NATIONAL BENTGRASS TEST (Fairway/Tee)

Entries and Sponsors

Entry No	Name	Species	Sponsor
1	DLFPS-AT/3026	Colonial	DLF Pickseed USA
*2	007	Creeping	Standard
*3	Pennncross	Creeping	Standard
*4	Crystal Blue Links	Creeping	Standard
*5	Greentime	Colonial	Standard
*6	L-93XD	Creeping	Jacklin Seed by Simplot®
*7	Armor	Creeping	Jacklin Seed by Simplot®
*8	Kingdom	Creeping	Jacklin Seed by Simplot®
*9	Nightlife	Creeping	Jacklin Seed by Simplot®
*10	V-8	Creeping	Jacklin Seed by Simplot®
*11	Piranha (DC-1)	Creeping	Mountain View Seeds & Toyo Green Co. Ltd.
*12	Musket (PPG-AT 104)	Colonial	Mountain View Seeds
*13	Chinook (H10G-OP)	Creeping	Vista Seed Partners LLC
*14	Shark	Creeping	Mountain View Seeds
*15	Barracuda	Creeping	Mountain View Seeds
16	PST-0RBS	Creeping	Pure Seed Testing Inc
*17	PC2.0 (PST-0CV6)	Creeping	Tee-2-Green

* Commercially Available in the USA in 2018

TABLE A. 2017 LOCATIONS, SITE DESCRIPTIONS AND MANAGEMENT PRACTICES IN THE 2014 NATIONAL BENTGRASS (FAIRWAY/TEE) TEST

LOCATION	SOIL TEXTURE	SOIL PH	SOIL PHOSPHOROUS (LBS/ACRE)	SOIL POTASSIUM (LBS/ACRE)	NITROGEN (LBS/1000 SQ FT)	SUN OR SHADE	MOWING HEIGHT (IN)	IRRIGATION PRACTICED
CA3	SANDY LOAM	7.1-7.5	0-60	241-375	2.1-3.0	FULL SUN	0.0-0.5	ONLY DURING SEVERE STRESS
IA1	SANDY CLAY LOAM	7.1-7.5	0-60	0-150	2.1-3.0	FULL SUN	0.0-0.5	TO PREVENT STRESS
IL1	SILT LOAM AND SILT	6.1-6.5	0-60	376-500	4.1-5.0	FULL SUN	0.0-0.5	TO PREVENT STRESS
IN1	SAND	7.6-8.5	61-150	151-240	3.1-4.0	FULL SUN	0.0-0.5	TO PREVENT DORMANCY
KS1	-	-	-	-	-	-	-	-
KY1	SILT LOAM AND SILT	4.6-5.5	451+	241-375	0.0-1.0	FULL SUN	0.6-1.0	TO PREVENT STRESS
MA1	LOAM	5.6-6.0	0-60	241-375	2.1-3.0	FULL SUN	0.0-0.5	TO PREVENT STRESS
MD1	SILT LOAM AND SILT	6.1-6.5	61-150	151-240	3.1-4.0	FULL SUN	0.0-0.5	TO PREVENT STRESS
MI1	-	-	-	-	-	-	-	-
MO1	-	-	-	-	-	-	-	-
ND1	SILTY CLAY AND CLAY	7.1-7.5	0-60	151-240	2.1-3.0	FULL SUN	0.0-0.5	TO PREVENT DORMANCY
NJ1	SANDY LOAM	-	-	-	1.1-2.0	FULL SUN	0.0-0.5	TO PREVENT STRESS
UT1	SILT LOAM AND SILT	7.1-7.5	0-60	0-150	-	FULL SUN	0.0-0.5	TO PREVENT DORMANCY
VA2	SILTY CLAY LOAM	6.1-6.5	61-150	151-240	3.1-4.0	FULL SUN	0.6-1.0	TO PREVENT DORMANCY
WA1	SILT LOAM AND SILT	5.6-6.0	451+	501+	2.1-3.0	FULL SUN	0.0-0.5	TO PREVENT STRESS

LOCATION	JANUARY QUALITY RATING	FEBRUARY QUALITY RATING	MARCH QUALITY RATING	APRIL QUALITY RATING	MAY QUALITY RATING	JUNE QUALITY RATING	JULY QUALITY RATING	AUGUST QUALITY RATING	SEPTEMBER QUALITY RATING	OCTOBER QUALITY RATING	NOVEMBER QUALITY RATING	DECEMBER QUALITY RATING	GENETIC COLOR	SPRING GREENUP	LEAF TEXTURE
CA3	X	X	X	X	X	X	X	X	X	X	X	X	X		
IA1				X	X	X	X	X	X	X			X		
IL1				X	X	X	X	X	X	X			X	X	
IN1					X	X	X	X	X	X	X		X		X
KS1				X	X	X	X	X	X	X					
KY1				X	X	X	X	X	X	X					
MA1			X	X	X	X	X	X	X	X				X	
MD1					X	X	X	X	X	X	X				
MI1					X	X	X	X	X	X			X	X	
MO1				X	X	X	X	X	X	X	X				
ND1					X	X	X	X	X	X					
NJ1				X	X	X	X	X	X	X				X	X
UT1				X	X	X	X	X	X	X				X	
VA2				X		X	X	X							
WA1					X	X	X	X	X	X			X		

TABLE B. (CONT'D)

LOCATIONS AND DATA COLLECTED IN 2017

LOCATION	SUMMER DENSITY	FALL DENSITY	DOLLAR SPOT	BROWN PATCH WARM TEMP.	ANTHRACNOSE RATINGS	FALL COLOR NOVEMBER	FALL COLOR DECEMBER	SEEDHEAD RATINGS	POA ANNUA	MOW QUALITY	LEAF RUST	PERCENT DOLLAR SPOT
CA3												
IA1			X	X								
IL1						X						
IN1	X									X		
KS1												
KY1												
MA1												X
MD1			X									
MI1						X						
MO1												
ND1	X	X										
NJ1		X	X		X	X						
UT1												
VA2												X
WA1												

TABLE 1A. MEAN TURFGRASS QUALITY RATINGS OF BENTGRASS CULTIVARS GROWN ON 1/
A FAIRWAY OR TEE GROWN IN LOCATION PERFORMANCE INDEX (LPI) GROUP 1 **/
2017 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/								
NAME	ENTRY #	NJ1	MD1	KY1	MI1	IA1	KS1	MEAN
* CHINOOK (H10G-OP)		7.3	7.1	7.9	6.7	6.5	6.4	7.0
* PIRANHA (DC-1)	13	7.3	7.7	7.6	6.4	5.9	5.8	6.8
* BARRACUDA	11	6.5	6.8	7.7	6.5	6.2	5.9	6.6
* SHARK	15	6.4	6.5	7.6	6.4	6.2	5.9	6.5
* 007	14	6.2	6.5	7.7	6.5	6.3	5.9	6.5
* V-8	2	6.2	6.1	7.7	6.5	6.4	6.0	6.5
* L-93XD	10	6.6	6.9	7.4	6.2	5.9	5.6	6.4
DLFPS-AT/3026	6	6.6	6.5	7.2	6.0	5.8	5.6	6.3
* CRYSTAL BLUE LINKS	1	5.4	5.4	7.8	6.6	6.6	6.0	6.3
* MUSKET (PPG-AT 104)	4	6.6	6.4	7.1	5.9	5.7	5.6	6.2
* NIGHTLIFE	12	4.7	5.3	7.7	6.6	6.5	5.7	6.1
* PC2.0 (PST-OCV6)	9	4.5	5.1	7.5	6.3	6.3	5.5	5.9
* GREENTIME	17	4.9	3.6	7.3	6.3	6.6	6.0	5.8
* KINGDOM	5	3.9	4.7	7.6	6.5	6.4	5.5	5.8
* ARMOR	8	3.8	4.3	7.5	6.4	6.5	5.5	5.7
PST-ORBS	7	4.0	4.4	7.3	6.2	6.3	5.4	5.6
* PENNCROSS	16	3.7	2.7	7.1	6.1	6.4	5.6	5.3
	3							
LSD VALUE		0.8	0.8	0.8	0.8	0.8	0.8	0.8
C.V. (%)		8.9	8.8	6.6	7.8	7.9	8.7	8.0

*/ COMMERCIALY AVAILABLE IN THE USA IN 2018

**/ ENTRIES WITHIN THIS TABLE ARE ORDERED BY THE OVERALL MEAN AND HAVE SIMILAR TURF QUALITY PERFORMANCES IN ALL TEST LOCATIONS INCLUDED IN THIS LPI GROUP. IF YOUR STATE IS NOT REPRESENTED, THEN CHOOSE A LPI GROUP THAT CONTAINS A LOCATION AND MANAGEMENT SIMILAR TO YOUR PLANTING CONDITIONS. FOR MORE INFORMATION ON LPI, GO TO WWW.NTEP.ORG/LPI_Q&A.PDF

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

Taken from the 2014 Bentgrass (Fairway/Tee) test - 2017 data report with permission from NTEP

TABLE 1B. MEAN TURFGRASS QUALITY RATINGS OF CREEPING BENTGRASS CULTIVARS GROWN ON 1/
A FAIRWAY OR TEE GROWN IN LOCATION PERFORMANCE INDEX (LPI) GROUP 1 */
2017 DATA
TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	NJ1	MD1	KY1	MI1	IA1	KS1	MEAN
CHINOOK (H10G-OP)	7.3	7.1	7.9	6.7	6.5	6.4	7.0
PIRANHA (DC-1)	7.3	7.7	7.6	6.4	5.9	5.8	6.8
BARRACUDA	6.5	6.8	7.7	6.5	6.2	5.9	6.6
SHARK	6.4	6.5	7.6	6.4	6.2	5.9	6.5
007	6.2	6.5	7.7	6.5	6.3	5.9	6.5
V-8	6.2	6.1	7.7	6.5	6.4	6.0	6.5
L-93XD	6.6	6.9	7.4	6.2	5.9	5.6	6.4
CRYSTAL BLUE LINKS	5.4	5.4	7.8	6.6	6.6	6.0	6.3
NIGHTLIFE	4.7	5.3	7.7	6.6	6.5	5.7	6.1
PC2.0 (PST-OCV6)	4.5	5.1	7.5	6.3	6.3	5.5	5.9
KINGDOM	3.9	4.7	7.6	6.5	6.4	5.5	5.8
ARMOR	3.8	4.3	7.5	6.4	6.5	5.5	5.7
PST-ORBS	4.0	4.4	7.3	6.2	6.3	5.4	5.6
PENNCROSS	3.7	2.7	7.1	6.1	6.4	5.6	5.3
LSD VALUE	0.8	0.8	0.8	0.8	0.8	0.8	0.8
C.V. (%)	9.1	8.8	6.6	7.7	7.9	8.7	8.0

TABLE 1C. MEAN TURFGRASS QUALITY RATINGS OF COLONIAL BENTGRASS CULTIVARS GROWN ON 1/
A FAIRWAY OR TEE GROWN IN LOCATION PERFORMANCE INDEX (LPI) GROUP 1 */
2017 DATA
TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	NJ1	MD1	KY1	MI1	IA1	KS1	MEAN
DLFPS-AT/3026	6.6	6.5	7.2	6.0	5.8	5.6	6.3
MUSKET (PPG-AT 104)	6.6	6.4	7.1	5.9	5.7	5.6	6.2
GREENTIME	4.9	3.6	7.3	6.3	6.6	6.0	5.8
LSD VALUE	0.8	0.8	0.8	0.8	0.8	0.8	0.8
C.V. (%)	8.2	9.0	6.9	8.2	8.3	8.7	8.2

*/ ENTRIES WITHIN THIS TABLE ARE ORDERED BY THE OVERALL MEAN AND HAVE SIMILAR TURF QUALITY PERFORMANCES IN ALL TEST LOCATIONS INCLUDED IN THIS LPI GROUP. IF YOUR STATE IS NOT REPRESENTED, THEN CHOOSE A LPI GROUP THAT CONTAINS A LOCATION AND MANAGEMENT SIMILAR TO YOUR PLANTING CONDITIONS. FOR MORE INFORMATION ON LPI, GO TO WWW.NTEP.ORG/LPI_Q&A.PDF

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 2A. MEAN TURFGRASS QUALITY RATINGS OF BENTGRASS CULTIVARS GROWN ON 1/
A FAIRWAY OR TEE GROWN IN LOCATION PERFORMANCE INDEX (LPI) GROUP 2 */
2017 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	IL1	MO1	WA1	MEAN
NIGHTLIFE	7.8	6.0	7.0	6.9
KINGDOM	7.8	6.0	7.1	6.9
ARMOR	7.7	5.6	6.8	6.7
PC2.0 (PST-OCV6)	7.5	5.7	6.6	6.6
CRYSTAL BLUE LINKS	7.7	5.5	6.4	6.6
007	7.4	5.8	6.3	6.6
BARRACUDA	7.3	5.8	6.2	6.6
PIRANHA (DC-1)	7.1	5.9	6.1	6.5
PST-ORBS	7.5	5.4	6.5	6.5
CHINOOK (H10G-OP)	7.4	5.4	5.8	6.4
V-8	7.4	5.4	6.0	6.4
SHARK	7.3	5.4	6.0	6.3
L-93XD	7.0	5.6	6.0	6.3
DLFPS-AT/3026	6.7	4.9	5.3	5.8
MUSKET (PPG-AT 104)	6.6	4.7	5.1	5.7
PENNCROSS	7.1	3.6	5.0	5.3
GREENTIME	7.1	3.6	4.8	5.3
LSD VALUE	0.8	0.8	0.8	0.8
C.V. (%)	6.8	9.4	8.2	7.9

*/ ENTRIES WITHIN THIS TABLE ARE ORDERED BY THE OVERALL MEAN AND HAVE SIMILAR TURF QUALITY PERFORMANCES IN ALL TEST LOCATIONS INCLUDED IN THIS LPI GROUP. IF YOUR STATE IS NOT REPRESENTED, THEN CHOOSE A LPI GROUP THAT CONTAINS A LOCATION AND MANAGEMENT SIMILAR TO YOUR PLANTING CONDITIONS. FOR MORE INFORMATION ON LPI, GO TO WWW.NTEP.ORG/LPI_Q&A.PDF

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 2B. MEAN TURFGRASS QUALITY RATINGS OF CREEPING BENTGRASS CULTIVARS GROWN ON 1/
A FAIRWAY OR TEE GROWN IN LOCATION PERFORMANCE INDEX (LPI) GROUP 2 */
2017 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/				
NAME	IL1	M01	WA1	MEAN
NIGHTLIFE	7.8	6.0	7.0	6.9
KINGDOM	7.8	6.0	7.1	6.9
ARMOR	7.7	5.6	6.8	6.7
PC2.0 (PST-OCV6)	7.5	5.7	6.6	6.6
CRYSTAL BLUE LINKS	7.7	5.5	6.4	6.6
007	7.4	5.8	6.3	6.6
BARRACUDA	7.3	5.8	6.2	6.6
PIRANHA (DC-1)	7.1	5.9	6.1	6.5
PST-ORBS	7.5	5.4	6.5	6.5
CHINOOK (H10G-OP)	7.4	5.4	5.8	6.4
V-8	7.4	5.4	6.0	6.4
SHARK	7.3	5.4	6.0	6.3
L-93XD	7.0	5.6	6.0	6.3
PENNCROSS	7.1	3.6	5.0	5.3
LSD VALUE	0.8	0.8	0.8	0.8
C.V. (%)	6.7	9.0	7.9	7.7

TABLE 2C. MEAN TURFGRASS QUALITY RATINGS OF COLONIAL BENTGRASS CULTIVARS GROWN ON 1/
A FAIRWAY OR TEE GROWN IN LOCATION PERFORMANCE INDEX (LPI) GROUP 2 */
2017 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/				
NAME	IL1	M01	WA1	MEAN
DLFPS-AT/3026	6.7	4.9	5.3	5.8
MUSKET (PPG-AT 104)	6.6	4.7	5.1	5.7
GREENTIME	7.1	3.6	4.8	5.3
LSD VALUE	0.8	0.8	0.8	0.8
C.V. (%)	7.3	11.3	9.9	8.9

*/ ENTRIES WITHIN THIS TABLE ARE ORDERED BY THE OVERALL MEAN AND HAVE SIMILAR TURF QUALITY PERFORMANCES IN ALL TEST LOCATIONS INCLUDED IN THIS LPI GROUP. IF YOUR STATE IS NOT REPRESENTED, THEN CHOOSE A LPI GROUP THAT CONTAINS A LOCATION AND MANAGEMENT SIMILAR TO YOUR PLANTING CONDITIONS. FOR MORE INFORMATION ON LPI, GO TO WWW.NTEP.ORG/LPI_Q&A.PDF

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3A.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER TRAFFIC STRESS AT AMHERST, MA 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	SPRING GREENUP	WEAR TOLERANCE RATINGS				QUALITY RATINGS							MEAN
		MAY	JULY	SEPTEMBER	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	
MUSKET (PPG-AT 104)	6.0	6.0	5.3	3.7	4.0	6.3	7.0	6.3	6.0	6.3	6.7	5.3	6.0
DLFPS-AT/3026	6.7	6.0	5.0	4.0	4.0	6.3	7.0	6.7	6.7	5.7	5.7	5.0	5.9
PIRANHA (DC-1)	4.7	6.7	5.3	7.0	2.3	5.7	6.0	6.3	6.7	6.0	6.7	6.7	5.8
KINGDOM	3.0	6.7	7.3	7.3	2.0	5.7	5.0	6.7	6.7	5.3	6.7	7.0	5.6
L-93XD	4.0	5.0	4.7	5.3	2.3	6.0	6.0	6.3	5.7	5.7	5.7	6.3	5.5
NIGHTLIFE	3.3	6.3	7.0	7.7	2.0	5.3	5.0	6.3	6.7	6.0	6.0	6.3	5.5
007	3.3	5.3	4.7	6.7	2.3	5.7	5.7	6.3	5.7	5.3	6.0	5.3	5.3
ARMOR	3.3	5.7	6.0	6.7	2.0	6.0	5.0	6.3	5.7	5.3	6.0	5.7	5.3
CHINOOK (H10G-OP)	5.0	5.3	4.7	6.0	2.3	5.3	6.0	6.7	5.0	5.7	6.0	5.7	5.3
V-8	2.7	5.0	4.7	5.7	2.0	5.0	5.0	5.7	5.7	5.3	6.7	6.0	5.2
GREENTIME	4.0	6.0	5.3	4.3	4.0	5.7	5.3	5.3	5.3	5.0	5.3	4.7	5.1
PC2.0 (PST-OCV6)	5.0	5.0	4.3	6.0	3.0	5.0	5.0	6.0	5.3	5.3	6.0	5.0	5.1
SHARK	3.0	4.7	4.7	7.0	2.0	5.0	5.7	6.3	5.3	5.3	5.7	5.7	5.1
BARRACUDA	4.0	4.7	5.0	6.0	2.7	5.0	5.3	5.3	5.3	5.0	5.3	5.3	4.9
PST-ORBS	4.0	4.0	4.0	6.0	2.3	5.3	5.0	6.0	5.0	5.0	5.3	5.0	4.9
CRYSTAL BLUE LINKS	3.0	5.0	4.0	5.3	2.3	4.7	4.3	5.0	5.0	4.7	5.0	4.7	4.5
PENNCROSS	2.7	3.7	3.7	3.7	1.7	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.7
LSD VALUE	0.9	0.9	0.6	0.8	0.6	0.8	0.4	0.9	0.7	0.8	0.7	0.7	0.4
C.V. (%)	14.6	10.8	8.1	9.3	16.0	8.4	5.5	8.6	7.7	8.9	7.8	7.8	4.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3B. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF CREEPING BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER TRAFFIC STRESS AT AMHERST, MA 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/													
NAME	SPRING GREENUP	WEAR TOLERANCE RATINGS			QUALITY RATINGS								MEAN
		MAY	JULY	SEPTEMBER	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	
PIRANHA (DC-1)	4.7	6.7	5.3	7.0	2.3	5.7	6.0	6.3	6.7	6.0	6.7	6.7	5.8
KINGDOM	3.0	6.7	7.3	7.3	2.0	5.7	5.0	6.7	6.7	5.3	6.7	7.0	5.6
L-93XD	4.0	5.0	4.7	5.3	2.3	6.0	6.0	6.3	5.7	5.7	5.7	6.3	5.5
NIGHTLIFE	3.3	6.3	7.0	7.7	2.0	5.3	5.0	6.3	6.7	6.0	6.0	6.3	5.5
007	3.3	5.3	4.7	6.7	2.3	5.7	5.7	6.3	5.7	5.3	6.0	5.3	5.3
ARMOR	3.3	5.7	6.0	6.7	2.0	6.0	5.0	6.3	5.7	5.3	6.0	5.7	5.3
CHINOOK (H10G-OP)	5.0	5.3	4.7	6.0	2.3	5.3	6.0	6.7	5.0	5.7	6.0	5.7	5.3
V-8	2.7	5.0	4.7	5.7	2.0	5.0	5.0	5.7	5.7	5.3	6.7	6.0	5.2
PC2.0 (PST-OCV6)	5.0	5.0	4.3	6.0	3.0	5.0	5.0	6.0	5.3	5.3	6.0	5.0	5.1
SHARK	3.0	4.7	4.7	7.0	2.0	5.0	5.7	6.3	5.3	5.3	5.7	5.7	5.1
BARRACUDA	4.0	4.7	5.0	6.0	2.7	5.0	5.3	5.3	5.3	5.0	5.3	5.3	4.9
PST-ORBS	4.0	4.0	4.0	6.0	2.3	5.3	5.0	6.0	5.0	5.0	5.3	5.0	4.9
CRYSTAL BLUE LINKS	3.0	5.0	4.0	5.3	2.3	4.7	4.3	5.0	5.0	4.7	5.0	4.7	4.5
PENNCROSS	2.7	3.7	3.7	3.7	1.7	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.7
LSD VALUE	1.1	0.8	0.6	0.9	1.1	0.7	0.5	0.8	0.7	0.9	0.7	0.6	0.3
C.V. (%)	17.3	9.7	7.8	9.0	20.1	7.8	5.6	8.4	8.2	9.0	7.2	7.2	4.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3C. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF COLONIAL BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER TRAFFIC STRESS AT AMHERST, MA 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/													
NAME	SPRING GREENUP	WEAR TOLERANCE RATINGS			QUALITY RATINGS								MEAN
		MAY	JULY	SEPTEMBER	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	
MUSKET (PPG-AT 104)	6.0	6.0	5.3	3.7	4	6.3	7.0	6.3	6.0	6.3	6.7	5.3	6.0
DLFPS-AT/3026	6.7	6.0	5.0	4.0	4	6.3	7.0	6.7	6.7	5.7	5.7	5.0	5.9
GREENTIME	4.0	6.0	5.3	4.3	4	5.7	5.3	5.3	5.3	5.0	5.3	4.7	5.1
LSD VALUE	0.8	1.3	1.2	0.9	0	1.5	0.8	1.2	0.9	1.3	1.5	1.3	0.7
C.V. (%)	6.0	9.6	10.1	10.2	0	10.9	5.2	8.6	6.8	10.2	11.3	11.5	5.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 4A.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER TRAFFIC STRESS AT BLACKSBURG, VA 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	DISEASE JULY	QUALITY RATINGS BEFORE TRAFFIC (APR-AUG)				QUALITY RATINGS FOR TRAFFIC				QUALITY RATINGS FOR DIVOT RECOVERY				MEAN
		APR	JUN	JUL	AUG	SEP	OCT_11	OCT_25	NOV	SEP	OCT_11	OCT_25	NOV	
GREENTIME	9.0	6.0	7.7	7.0	7.3	6.0	6.7	7.3	7.0	7.3	7.7	8.0	7.7	7.1
CRYSTAL BLUE LINKS	7.0	6.7	6.7	6.0	7.7	6.3	6.0	7.3	7.0	6.7	7.7	7.7	7.3	6.9
DLFPS-AT/3026	7.0	7.0	6.3	6.7	7.3	4.3	6.0	7.0	6.3	7.7	8.0	8.0	8.0	6.9
PENNCROSS	8.7	6.0	7.3	6.7	7.3	5.3	5.3	6.7	7.3	7.3	7.3	7.7	7.3	6.8
MUSKET (PPG-AT 104)	8.0	7.0	6.3	6.0	7.7	5.3	6.0	7.0	6.7	7.0	6.7	7.3	7.3	6.7
V-8	7.0	5.7	6.3	6.0	7.3	5.7	5.7	7.0	7.0	7.0	7.0	8.0	7.3	6.7
PST-ORBS	6.3	6.3	6.0	6.0	7.3	5.3	5.7	7.0	7.0	7.0	7.0	7.3	7.0	6.6
SHARK	5.7	7.0	5.3	5.0	7.0	5.3	6.0	7.0	7.0	7.0	7.0	8.0	7.3	6.6
ARMOR	6.7	6.0	6.3	5.7	6.3	5.3	5.7	7.0	7.0	6.7	6.7	7.7	7.0	6.4
KINGDOM	6.7	6.0	6.0	5.3	6.0	5.3	6.0	7.3	7.0	6.7	7.0	7.7	7.0	6.4
NIGHTLIFE	5.3	6.3	5.7	5.0	5.7	5.7	6.0	7.3	7.0	6.7	6.7	7.7	7.0	6.4
PC2.0 (PST-OCV6)	7.3	6.0	6.3	5.3	6.7	4.7	5.7	7.0	7.0	6.7	7.0	7.7	7.0	6.4
007	6.3	6.3	5.7	5.0	6.7	5.0	5.0	6.7	6.7	7.0	7.3	7.3	7.3	6.3
BARRACUDA	7.0	6.0	6.3	5.7	6.3	4.7	5.3	6.7	6.7	6.7	6.7	7.3	7.3	6.3
L-93XD	6.7	6.0	6.0	5.3	6.7	5.0	5.3	6.7	6.3	6.7	6.7	7.3	7.3	6.3
PIRANHA (DC-1)	6.3	7.0	5.3	5.3	7.0	4.7	4.7	6.7	6.3	6.3	6.7	7.3	7.3	6.2
CHINOOK (H10G-OP)	5.3	6.7	5.3	4.0	5.7	5.3	5.7	7.0	6.7	6.7	6.3	7.3	7.0	6.1
LSD VALUE	2.5	0.8	1.1	1.3	1.6	1.5	1.6	2.0	0.9	1.3	0.8	1.3	1.2	0.4
C.V. (%)	17.3	6.6	9.6	12.5	11.1	12.5	11.3	8.4	5.7	7.2	6.0	6.0	5.9	3.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 4B.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF CREEPING BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER TRAFFIC STRESS AT BLACKSBURG, VA 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	DISEASE JULY	QUALITY RATINGS BEFORE TRAFFIC (APR-AUG)				QUALITY RATINGS FOR TRAFFIC				QUALITY RATINGS FOR DIVOT RECOVERY				MEAN
		APR	JUN	JUL	AUG	SEP	OCT_11	OCT_25	NOV	SEP	OCT_11	OCT_25	NOV	
CRYSTAL BLUE LINKS	7.0	6.7	6.7	6.0	7.7	6.3	6.0	7.3	7.0	6.7	7.7	7.7	7.3	6.9
PENNCROSS	8.7	6.0	7.3	6.7	7.3	5.3	5.3	6.7	7.3	7.3	7.3	7.7	7.3	6.8
V-8	7.0	5.7	6.3	6.0	7.3	5.7	5.7	7.0	7.0	7.0	7.0	8.0	7.3	6.7
PST-ORBS	6.3	6.3	6.0	6.0	7.3	5.3	5.7	7.0	7.0	7.0	7.0	7.3	7.0	6.6
SHARK	5.7	7.0	5.3	5.0	7.0	5.3	6.0	7.0	7.0	7.0	7.0	8.0	7.3	6.6
ARMOR	6.7	6.0	6.3	5.7	6.3	5.3	5.7	7.0	7.0	6.7	6.7	7.7	7.0	6.4
KINGDOM	6.7	6.0	6.0	5.3	6.0	5.3	6.0	7.3	7.0	6.7	7.0	7.7	7.0	6.4
NIGHTLIFE	5.3	6.3	5.7	5.0	5.7	5.7	6.0	7.3	7.0	6.7	6.7	7.7	7.0	6.4
PC2.0 (PST-OCV6)	7.3	6.0	6.3	5.3	6.7	4.7	5.7	7.0	7.0	6.7	7.0	7.7	7.0	6.4
007	6.3	6.3	5.7	5.0	6.7	5.0	5.0	6.7	6.7	7.0	7.3	7.3	7.3	6.3
BARRACUDA	7.0	6.0	6.3	5.7	6.3	4.7	5.3	6.7	6.7	6.7	6.7	7.3	7.3	6.3
L-93XD	6.7	6.0	6.0	5.3	6.7	5.0	5.3	6.7	6.3	6.7	6.7	7.3	7.3	6.3
PIRANHA (DC-1)	6.3	7.0	5.3	5.3	7.0	4.7	4.7	6.7	6.3	6.3	6.7	7.3	7.3	6.2
CHINOOK (H10G-OP)	5.3	6.7	5.3	4.0	5.7	5.3	5.7	7.0	6.7	6.7	6.3	7.3	7.0	6.1
LSD VALUE	3.0	1.0	1.2	1.4	1.8	1.5	1.7	1.8	0.9	1.4	0.8	1.2	1.3	0.6
C.V. (%)	18.5	7.4	9.9	12.6	11.9	11.8	11.5	8.2	5.6	7.1	5.8	5.8	5.8	4.1

TABLE 4C.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF COLONIAL BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER TRAFFIC STRESS AT BLACKSBURG, VA 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	DISEASE JULY	QUALITY RATINGS BEFORE TRAFFIC (APR-AUG)				QUALITY RATINGS FOR TRAFFIC				QUALITY RATINGS FOR DIVOT RECOVERY				MEAN
		APR	JUN	JUL	AUG	SEP	OCT_11	OCT_25	NOV	SEP	OCT_11	OCT_25	NOV	
GREENTIME	9.0	6	7.7	7.0	7.3	6.0	6.7	7.3	7.0	7.3	7.7	8.0	7.7	7.1
DLFPS-AT/3026	7.0	7	6.3	6.7	7.3	4.3	6.0	7.0	6.3	7.7	8.0	8.0	8.0	6.9
MUSKET (PPG-AT 104)	8.0	7	6.3	6.0	7.7	5.3	6.0	7.0	6.7	7.0	6.7	7.3	7.3	6.7
LSD VALUE	2.1	0	1.5	1.5	1.5	2.0	1.5	1.8	0.9	1.3	1.2	0.8	0.9	0.3
C.V. (%)	11.4	0	9.8	10.2	9.0	16.9	10.7	11.0	6.1	7.9	7.1	4.3	5.3	2.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5A.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER REDUCED IRRIGATION AT RIVERSIDE, CA 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	GENETIC COLOR	QUALITY RATINGS												MEAN
		JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	
007	7.0	6.3	6.0	6.7	7.3	6.7	6.3	6.3	6.0	6.0	6.3	6.0	6.3	6.4
L-93XD	7.0	6.0	5.7	6.0	6.7	6.0	6.3	6.7	6.3	6.7	6.7	6.0	6.0	6.3
PENNCROSS	7.0	6.3	6.3	6.7	6.7	6.7	6.0	6.3	6.0	6.0	6.0	6.3	6.0	6.3
PST-ORBS	7.0	6.3	6.0	6.0	7.0	6.7	6.3	6.7	6.0	6.3	6.0	5.7	6.3	6.3
SHARK	7.0	6.3	6.0	6.3	6.3	6.7	6.3	6.0	6.0	6.7	6.0	6.3	6.3	6.3
CHINOOK (H10G-OP)	7.3	6.0	5.7	6.0	6.0	6.7	6.0	7.0	6.3	6.7	6.0	6.0	6.0	6.2
CRYSTAL BLUE LINKS	7.7	6.0	6.0	6.3	5.7	6.3	6.3	6.3	6.3	6.3	6.3	6.0	6.0	6.2
PC2.0 (PST-OCV6)	7.7	6.0	6.0	6.3	6.3	7.0	6.0	6.3	6.0	6.0	6.3	6.0	6.0	6.2
PIRANHA (DC-1)	8.0	6.0	5.7	6.0	6.7	6.7	6.3	6.3	6.3	6.3	6.3	6.0	6.0	6.2
BARRACUDA	7.3	6.0	6.0	6.3	6.3	6.3	5.7	6.0	6.3	6.3	6.0	6.0	6.0	6.1
KINGDOM	8.0	5.7	6.0	6.0	6.7	6.3	6.0	6.3	5.7	6.0	6.0	6.0	6.0	6.1
NIGHTLIFE	8.0	5.7	6.0	6.0	6.3	6.7	6.3	6.3	6.0	6.3	6.3	5.7	5.0	6.1
ARMOR	8.0	6.0	5.7	5.7	6.0	6.0	5.7	6.3	6.3	6.3	5.7	6.0	6.0	6.0
DLFPS-AT/3026	7.0	6.0	5.7	5.7	6.0	6.0	5.7	6.3	6.0	5.7	5.3	6.0	6.0	5.9
MUSKET (PPG-AT 104)	7.0	6.0	5.7	5.7	6.0	6.0	5.7	6.0	6.0	6.0	6.0	6.0	6.0	5.9
V-8	7.0	5.7	5.7	5.7	6.7	6.7	5.7	6.0	6.0	6.0	5.7	6.0	5.7	5.9
GREENTIME	7.3	5.3	5.3	6.0	6.0	6.0	5.7	6.3	5.7	6.3	5.3	5.7	6.0	5.8
LSD VALUE	0.5	1.0	1.3	1.0	1.2	1.1	2.0	1.8	2.0	2.1	1.2	1.3	0.5	0.5
C.V. (%)	4.2	6.7	7.4	7.2	8.4	7.1	10.2	8.8	9.3	10.2	8.4	6.8	4.7	3.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5B. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF CREEPING BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER REDUCED IRRIGATION AT RIVERSIDE, CA 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/														
NAME	GENETIC COLOR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
007	7.0	6.3	6.0	6.7	7.3	6.7	6.3	6.3	6.0	6.0	6.3	6.0	6.3	6.4
L-93XD	7.0	6.0	5.7	6.0	6.7	6.0	6.3	6.7	6.3	6.7	6.7	6.0	6.0	6.3
PENNCROSS	7.0	6.3	6.3	6.7	6.7	6.7	6.0	6.3	6.0	6.0	6.0	6.3	6.0	6.3
PST-ORBS	7.0	6.3	6.0	6.0	7.0	6.7	6.3	6.7	6.0	6.3	6.0	5.7	6.3	6.3
SHARK	7.0	6.3	6.0	6.3	6.3	6.7	6.3	6.0	6.0	6.7	6.0	6.3	6.3	6.3
CHINOOK (H10G-OP)	7.3	6.0	5.7	6.0	6.0	6.7	6.0	7.0	6.3	6.7	6.0	6.0	6.0	6.2
CRYSTAL BLUE LINKS	7.7	6.0	6.0	6.3	5.7	6.3	6.3	6.3	6.3	6.3	6.3	6.0	6.0	6.2
PC2.0 (PST-OCV6)	7.7	6.0	6.0	6.3	6.3	7.0	6.0	6.3	6.0	6.0	6.3	6.0	6.0	6.2
PIRANHA (DC-1)	8.0	6.0	5.7	6.0	6.7	6.7	6.3	6.3	6.3	6.3	6.3	6.0	6.0	6.2
BARRACUDA	7.3	6.0	6.0	6.3	6.3	6.3	5.7	6.0	6.3	6.3	6.0	6.0	6.0	6.1
KINGDOM	8.0	5.7	6.0	6.0	6.7	6.3	6.0	6.3	5.7	6.0	6.0	6.0	6.0	6.1
NIGHTLIFE	8.0	5.7	6.0	6.0	6.3	6.7	6.3	6.3	6.0	6.3	6.3	5.7	5.0	6.1
ARMOR	8.0	6.0	5.7	5.7	6.0	6.0	5.7	6.3	6.3	6.3	5.7	6.0	6.0	6.0
V-8	7.0	5.7	5.7	5.7	6.7	6.7	5.7	6.0	6.0	6.0	5.7	6.0	5.7	5.9
LSD VALUE	0.5	1.1	1.1	1.1	1.2	1.4	1.8	1.5	2.1	2.2	1.3	1.3	0.6	0.6
C.V. (%)	3.8	6.8	6.6	7.3	8.2	7.6	9.6	8.1	10.0	10.5	7.8	7.0	5.2	3.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5C. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF COLONIAL BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER REDUCED IRRIGATION AT RIVERSIDE, CA 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/														
NAME	GENETIC COLOR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
DLFPS-AT/3026	7.0	6.0	5.7	5.7	6.0	6	5.7	6.3	6.0	5.7	5.3	6.0	6	5.9
MUSKET (PPG-AT 104)	7.0	6.0	5.7	5.7	6.0	6	5.7	6.0	6.0	6.0	6.0	6.0	6	5.9
GREENTIME	7.3	5.3	5.3	6.0	6.0	6	5.7	6.3	5.7	6.3	5.3	5.7	6	5.8
LSD VALUE	0.8	0.8	1.5	0.8	1.3	0	0.0	1.8	0.8	0.9	0.8	0.8	0	0.4
C.V. (%)	4.7	5.8	12.0	5.8	9.6	0	0.0	12.6	5.7	6.8	6.0	5.7	0	2.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6A.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER REDUCED IRRIGATION AT WEST LAFAYETTE, IN 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	GENETIC COLOR	LEAF TEXTURE	SUMMER DENSITY	MOW QUALITY	MAY	JUN	QUALITY RATINGS					
							JUL	AUG	SEP	OCT	NOV	MEAN
NIGHTLIFE	8.0	7.7	8.0	7.3	7.3	8.0	8.3	8.0	7.0	7.7	6.0	7.5
PC2.0 (PST-OCV6)	6.7	7.7	7.7	7.0	7.7	8.3	7.7	7.3	7.0	7.0	6.0	7.3
ARMOR	8.3	7.3	8.7	6.0	6.3	7.3	7.7	7.7	7.0	7.3	7.3	7.2
CHINOOK (H10G-OP)	7.0	6.7	8.3	6.3	6.7	7.0	7.0	7.3	6.3	7.3	5.7	6.8
V-8	6.7	7.0	8.0	6.7	6.0	6.7	7.0	7.7	7.0	7.3	6.0	6.8
BARRACUDA	7.0	7.0	7.3	6.7	5.7	6.7	7.0	7.7	6.7	6.7	6.3	6.7
L-93XD	6.7	7.0	8.0	5.7	6.0	7.0	7.0	7.7	6.3	6.7	6.0	6.7
PST-ORBS	7.3	7.7	8.7	4.3	5.7	6.7	7.3	7.3	6.0	7.3	5.7	6.6
CRYSTAL BLUE LINKS	6.3	6.7	7.3	6.7	6.3	6.3	6.7	7.3	6.7	6.7	5.3	6.5
KINGDOM	8.3	7.3	8.0	5.0	5.7	6.7	7.0	7.0	6.3	7.0	6.0	6.5
PIRANHA (DC-1)	7.0	7.7	8.0	5.0	5.7	7.0	7.0	6.7	7.0	6.7	5.3	6.5
007	7.0	7.3	9.0	4.7	5.0	6.7	7.0	7.0	6.3	7.3	5.7	6.4
SHARK	6.7	7.0	8.0	5.3	6.0	6.7	7.0	7.0	6.0	6.3	6.0	6.4
MUSKET (PPG-AT 104)	5.3	7.3	7.7	6.7	6.0	6.7	6.0	6.3	6.7	5.7	6.3	6.2
DLFPS-AT/3026	5.3	6.3	7.3	8.0	5.7	6.0	6.3	6.7	6.3	5.3	4.7	5.9
PENNCROSS	6.7	5.0	5.7	8.3	5.7	6.3	6.3	6.3	5.7	5.7	5.0	5.9
GREENTIME	7.7	6.7	7.0	7.3	5.3	5.3	5.7	6.3	5.0	5.7	4.3	5.4
LSD VALUE	0.7	1.2	0.8	2.0	2.3	2.0	1.0	1.4	1.6	1.0	1.6	0.7
C.V. (%)	6.1	9.4	6.6	17.2	15.7	12.9	8.3	8.9	10.8	9.0	13.6	6.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6B. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF CREEPING BENTGRASS CULTIVARS GROWN ON A FAIRWAY OR TEE UNDER REDUCED IRRIGATION AT WEST LAFAYETTE, IN 1/ 2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/												
NAME	GENETIC COLOR	LEAF TEXTURE	SUMMER DENSITY	MOW QUALITY	MAY	JUN	QUALITY RATINGS					MEAN
							JUL	AUG	SEP	OCT	NOV	
NIGHTLIFE	8.0	7.7	8.0	7.3	7.3	8.0	8.3	8.0	7.0	7.7	6.0	7.5
PC2.0 (PST-OCV6)	6.7	7.7	7.7	7.0	7.7	8.3	7.7	7.3	7.0	7.0	6.0	7.3
ARMOR	8.3	7.3	8.7	6.0	6.3	7.3	7.7	7.7	7.0	7.3	7.3	7.2
CHINOOK (H10G-OP)	7.0	6.7	8.3	6.3	6.7	7.0	7.0	7.3	6.3	7.3	5.7	6.8
V-8	6.7	7.0	8.0	6.7	6.0	6.7	7.0	7.7	7.0	7.3	6.0	6.8
BARRACUDA	7.0	7.0	7.3	6.7	5.7	6.7	7.0	7.7	6.7	6.7	6.3	6.7
L-93XD	6.7	7.0	8.0	5.7	6.0	7.0	7.0	7.7	6.3	6.7	6.0	6.7
PST-ORBS	7.3	7.7	8.7	4.3	5.7	6.7	7.3	7.3	6.0	7.3	5.7	6.6
CRYSTAL BLUE LINKS	6.3	6.7	7.3	6.7	6.3	6.3	6.7	7.3	6.7	6.7	5.3	6.5
KINGDOM	8.3	7.3	8.0	5.0	5.7	6.7	7.0	7.0	6.3	7.0	6.0	6.5
PIRANHA (DC-1)	7.0	7.7	8.0	5.0	5.7	7.0	7.0	6.7	7.0	6.7	5.3	6.5
007	7.0	7.3	9.0	4.7	5.0	6.7	7.0	7.0	6.3	7.3	5.7	6.4
SHARK	6.7	7.0	8.0	5.3	6.0	6.7	7.0	7.0	6.0	6.3	6.0	6.4
PENNCROSS	6.7	5.0	5.7	8.3	5.7	6.3	6.3	6.3	5.7	5.7	5.0	5.9
LSD VALUE	0.7	1.1	0.7	2.0	2.5	2.2	1.3	1.2	2.0	1.2	2.0	0.8
C.V. (%)	6.0	8.6	5.7	18.0	16.8	12.4	8.2	7.6	11.4	8.4	13.9	6.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6C. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF COLONIAL BENTGRASS CULTIVARS GROWN ON A FAIRWAY OR TEE UNDER REDUCED IRRIGATION AT WEST LAFAYETTE, IN 1/ 2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/												
NAME	GENETIC COLOR	LEAF TEXTURE	SUMMER DENSITY	MOW QUALITY	MAY	JUN	QUALITY RATINGS					MEAN
							JUL	AUG	SEP	OCT	NOV	
MUSKET (PPG-AT 104)	5.3	7.3	7.7	6.7	6.0	6.7	6.0	6.3	6.7	5.7	6.3	6.2
DLFPS-AT/3026	5.3	6.3	7.3	8.0	5.7	6.0	6.3	6.7	6.3	5.3	4.7	5.9
GREENTIME	7.7	6.7	7.0	7.3	5.3	5.3	5.7	6.3	5.0	5.7	4.3	5.4
LSD VALUE	0.8	2.0	1.9	1.9	0.9	1.9	1.3	2.0	0.9	1.8	1.5	0.8
C.V. (%)	5.5	13.0	11.1	11.1	7.2	13.6	9.6	13.7	6.8	14.1	13.0	5.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7A. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER REDUCED IRRIGATION AT LOGAN, UT 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	SPRING GREENUP	APR	MAY	JUN	QUALITY RATINGS			SEP	OCT	MEAN
					JUL	AUG				
MUSKET (PPG-AT 104)	7.0	5.0	7.3	6.0	7.0	6.7		5.7	5.7	6.2
NIGHTLIFE	6.7	6.0	8.7	4.0	6.0	6.0		6.0	6.0	6.1
DLFPS-AT/3026	7.0	5.7	7.3	6.0	6.7	6.0		5.3	4.7	6.0
ARMOR	7.3	5.3	8.0	4.0	6.0	6.0		6.3	5.3	5.9
KINGDOM	6.3	5.7	9.0	5.0	6.0	5.0		5.7	4.7	5.9
CRYSTAL BLUE LINKS	6.7	6.0	7.0	3.7	6.0	5.3		5.0	6.0	5.6
PIRANHA (DC-1)	6.7	6.3	7.7	3.7	6.0	4.7		5.0	5.3	5.5
SHARK	6.7	5.7	6.7	3.0	4.7	6.0		5.3	5.7	5.3
CHINOOK (H10G-OP)	7.0	5.7	7.3	3.3	5.0	5.0		5.0	5.0	5.2
V-8	6.0	6.0	7.3	3.3	5.7	4.7		4.3	5.3	5.2
BARRACUDA	7.0	5.3	7.3	3.3	5.7	5.3		4.0	4.7	5.1
PC2.0 (PST-OCV6)	6.7	5.7	7.3	3.3	5.3	4.7		4.0	4.7	5.0
PST-ORBS	6.3	5.0	7.0	3.0	4.7	5.0		4.7	5.3	5.0
007	6.3	6.0	6.7	2.3	4.3	4.7		4.7	4.7	4.8
PENNCROSS	5.7	4.3	5.0	2.0	5.3	5.3		5.3	5.0	4.6
GREENTIME	6.3	5.0	5.0	2.0	3.7	4.7		4.7	4.7	4.2
L-93XD	7.0	5.3	7.0	3.0	3.7	3.7		3.7	3.3	4.2
LSD VALUE	1.0	1.1	1.0	0.9	1.5	2.6		1.9	2.9	0.9
C.V. (%)	7.5	10.0	8.4	17.0	15.7	20.1		18.0	20.9	9.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7B. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF CREEPING BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER REDUCED IRRIGATION AT LOGAN, UT 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	SPRING GREENUP	APR	MAY	JUN	QUALITY RATINGS			SEP	OCT	MEAN
NIGHTLIFE	6.7	6.0	8.7	4.0	6.0	6.0	6.0	6.0	6.0	6.1
ARMOR	7.3	5.3	8.0	4.0	6.0	6.0	6.3	5.3	5.3	5.9
KINGDOM	6.3	5.7	9.0	5.0	6.0	5.0	5.7	4.7	5.7	5.9
CRYSTAL BLUE LINKS	6.7	6.0	7.0	3.7	6.0	5.3	5.0	6.0	6.0	5.6
PIRANHA (DC-1)	6.7	6.3	7.7	3.7	6.0	4.7	5.0	5.3	5.3	5.5
SHARK	6.7	5.7	6.7	3.0	4.7	6.0	5.3	5.7	5.7	5.3
CHINOOK (H10G-OP)	7.0	5.7	7.3	3.3	5.0	5.0	5.0	5.0	5.0	5.2
V-8	6.0	6.0	7.3	3.3	5.7	4.7	4.3	5.3	5.3	5.2
BARRACUDA	7.0	5.3	7.3	3.3	5.7	5.3	4.0	4.7	4.7	5.1
PC2.0 (PST-OCV6)	6.7	5.7	7.3	3.3	5.3	4.7	4.0	4.7	4.7	5.0
PST-ORBS	6.3	5.0	7.0	3.0	4.7	5.0	4.7	5.3	5.3	5.0
007	6.3	6.0	6.7	2.3	4.3	4.7	4.7	4.7	4.7	4.8
PENNCROSS	5.7	4.3	5.0	2.0	5.3	5.3	5.3	5.0	5.0	4.6
L-93XD	7.0	5.3	7.0	3.0	3.7	3.7	3.7	3.3	3.3	4.2
LSD VALUE	1.0	1.1	0.7	1.2	1.9	3.1	1.7	2.4	2.4	1.1
C.V. (%)	7.5	9.8	5.9	20.0	16.9	21.8	17.0	19.4	19.4	10.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7C. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF COLONIAL BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER REDUCED IRRIGATION AT LOGAN, UT 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	SPRING GREENUP	APR	MAY	JUN	QUALITY RATINGS			SEP	OCT	MEAN
MUSKET (PPG-AT 104)	7.0	5.0	7.3	6	7.0	6.7	5.7	5.7	5.7	6.2
DLFPS-AT/3026	7.0	5.7	7.3	6	6.7	6.0	5.3	4.7	4.7	6.0
GREENTIME	6.3	5.0	5.0	2	3.7	4.7	4.7	4.7	4.7	4.2
LSD VALUE	0.8	1.2	2.8	0	1.2	1.5	2.3	2.3	2.3	0.8
C.V. (%)	4.9	10.1	18.3	0	9.1	11.5	18.6	20.0	20.0	5.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 8A.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER LOW TEMPERATURE/WINTER TOLERANCE AT FARGO, ND 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	DENSITY RATINGS			PERCENT GROUND COVER			QUALITY RATINGS					MEAN
	SPRING	SUMMER	FALL	SPRING	SUMMER	FALL	MAY	JUN	JUL	AUG	SEP	
KINGDOM	5.0	6	7.0	93.3	93.3	99.0	5.7	6.0	6.7	7.0	7.3	6.5
SHARK	5.0	6	7.0	95.0	99.0	99.0	6.7	6.7	6.0	7.3	6.0	6.5
PST-ORBS	4.0	6	7.0	95.0	95.0	99.0	5.7	6.0	6.0	7.0	7.0	6.3
CRYSTAL BLUE LINKS	6.0	6	7.0	90.0	95.0	99.0	6.0	6.0	6.0	7.0	6.0	6.2
PIRANHA (DC-1)	5.0	6	6.0	93.3	95.0	99.0	6.0	6.0	6.0	7.0	6.0	6.2
007	4.0	6	6.0	95.0	95.0	96.3	5.7	5.7	6.0	6.7	6.3	6.1
CHINOOK (H10G-OP)	5.0	6	7.0	90.0	95.0	96.0	6.3	6.0	6.0	6.3	6.0	6.1
L-93XD	5.0	6	7.0	85.0	90.0	95.0	5.3	5.3	6.0	7.0	7.0	6.1
MUSKET (PPG-AT 104)	4.0	5	7.0	85.0	95.0	99.0	4.7	5.3	6.0	7.0	7.7	6.1
PC2.0 (PST-OCV6)	4.0	6	6.3	95.0	99.0	99.0	5.3	5.3	6.7	7.0	6.3	6.1
PENNCROSS	5.3	6	6.0	95.0	90.0	99.0	4.7	5.7	7.0	7.0	6.0	6.1
ARMOR	5.0	6	6.0	83.3	90.0	95.0	5.3	5.3	6.3	6.0	6.3	5.9
BARRACUDA	5.0	6	7.0	93.3	99.0	99.0	5.7	5.7	5.0	7.0	6.0	5.9
DLFPS-AT/3026	4.0	6	6.0	85.0	95.0	95.0	4.7	5.7	6.3	6.3	6.3	5.9
NIGHTLIFE	4.0	5	7.0	88.3	95.0	99.0	5.0	5.3	6.0	6.0	7.0	5.9
V-8	4.0	6	7.0	93.3	95.0	99.0	5.7	5.7	5.0	6.7	6.0	5.8
GREENTIME	4.7	5	7.0	83.3	91.7	95.0	4.7	5.0	6.0	6.3	6.7	5.7
LSD VALUE	0.3	0	0.2	4.6	1.5	2.3	0.8	0.9	0.4	0.6	0.6	0.4
C.V. (%)	4.4	0	2.1	3.2	1.1	1.4	9.0	7.8	4.7	4.9	5.8	3.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

Taken from the 2014 Bentgrass (Fairway/Tee) test - 2017 data report with permission from NTEP

TABLE 8B.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF CREEPING BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER LOW TEMPERATURE/WINTER TOLERANCE AT FARGO, ND 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	DENSITY RATINGS			PERCENT GROUND COVER			QUALITY RATINGS					MEAN
	SPRING	SUMMER	FALL	SPRING	SUMMER	FALL	MAY	JUN	JUL	AUG	SEP	
KINGDOM	5.0	6	7.0	93.3	93.3	99.0	5.7	6.0	6.7	7.0	7.3	6.5
SHARK	5.0	6	7.0	95.0	99.0	99.0	6.7	6.7	6.0	7.3	6.0	6.5
PST-ORBS	4.0	6	7.0	95.0	95.0	99.0	5.7	6.0	6.0	7.0	7.0	6.3
CRYSTAL BLUE LINKS	6.0	6	7.0	90.0	95.0	99.0	6.0	6.0	6.0	7.0	6.0	6.2
PIRANHA (DC-1)	5.0	6	6.0	93.3	95.0	99.0	6.0	6.0	6.0	7.0	6.0	6.2
007	4.0	6	6.0	95.0	95.0	96.3	5.7	5.7	6.0	6.7	6.3	6.1
CHINOOK (H10G-OP)	5.0	6	7.0	90.0	95.0	96.0	6.3	6.0	6.0	6.3	6.0	6.1
L-93XD	5.0	6	7.0	85.0	90.0	95.0	5.3	5.3	6.0	7.0	7.0	6.1
PC2.0 (PST-OCV6)	4.0	6	6.3	95.0	99.0	99.0	5.3	5.3	6.7	7.0	6.3	6.1
PENNCROSS	5.3	6	6.0	95.0	90.0	99.0	4.7	5.7	7.0	7.0	6.0	6.1
ARMOR	5.0	6	6.0	83.3	90.0	95.0	5.3	5.3	6.3	6.0	6.3	5.9
BARRACUDA	5.0	6	7.0	93.3	99.0	99.0	5.7	5.7	5.0	7.0	6.0	5.9
NIGHTLIFE	4.0	5	7.0	88.3	95.0	99.0	5.0	5.3	6.0	6.0	7.0	5.9
V-8	4.0	6	7.0	93.3	95.0	99.0	5.7	5.7	5.0	6.7	6.0	5.8
LSD VALUE	0.2	0	0.2	5.3	1.2	2.7	0.8	0.9	0.4	0.5	0.5	0.3
C.V. (%)	3.3	0	2.3	3.4	0.8	1.5	8.2	7.4	4.5	4.3	4.7	3.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 8C.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF COLONIAL BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE UNDER LOW TEMPERATURE/WINTER TOLERANCE AT FARGO, ND 1/
2017 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

NAME	DENSITY RATINGS			PERCENT GROUND COVER			QUALITY RATINGS					MEAN
	SPRING	SUMMER	FALL	SPRING	SUMMER	FALL	MAY	JUN	JUL	AUG	SEP	
MUSKET (PPG-AT 104)	4.0	5	7	85.0	95.0	99	4.7	5.3	6.0	7.0	7.7	6.1
DLFPS-AT/3026	4.0	6	6	85.0	95.0	95	4.7	5.7	6.3	6.3	6.3	5.9
GREENTIME	4.7	5	7	83.3	91.7	95	4.7	5.0	6.0	6.3	6.7	5.7
LSD VALUE	0.8	0	0	3.9	3.9	0	1.3	1.3	0.8	1.2	1.5	0.8
C.V. (%)	7.9	0	0	2.0	1.8	0	12.4	10.8	5.5	8.0	9.7	5.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9A. GENETIC COLOR RATINGS OF BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME	IA1	IL1	MI1	WA1	MEAN
KINGDOM	9.0	7.7	7.7	8.0	8.1
NIGHTLIFE	8.3	8.0	7.7	7.7	7.9
ARMOR	8.0	7.0	7.3	8.0	7.6
PIRANHA (DC-1)	8.3	7.0	7.0	6.0	7.1
CHINOOK (H10G-OP)	8.0	7.7	7.3	5.3	7.1
007	7.3	7.3	7.0	6.0	6.9
V-8	7.7	7.3	7.3	5.3	6.9
CRYSTAL BLUE LINKS	6.7	7.7	7.0	5.7	6.8
PST-ORBS	7.3	7.7	7.0	5.0	6.8
SHARK	7.0	7.7	7.0	5.3	6.8
PC2.0 (PST-OCV6)	7.3	7.0	7.0	5.3	6.7
GREENTIME	7.0	6.7	6.7	6.0	6.6
PENNCROSS	7.0	7.0	7.0	5.3	6.6
BARRACUDA	6.3	7.7	7.0	5.0	6.5
L-93XD	6.0	7.3	7.0	5.0	6.3
MUSKET (PPG-AT 104)	7.0	6.0	6.3	6.0	6.3
DLFPS-AT/3026	6.3	6.0	6.0	6.0	6.1
LSD VALUE	1.2	0.7	0.6	0.7	0.4
C.V. (%)	10.5	6.1	5.3	7.5	7.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9B. GENETIC COLOR RATINGS OF CREEPING BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME	IA1	IL1	MI1	WA1	MEAN
KINGDOM	9.0	7.7	7.7	8.0	8.1
NIGHTLIFE	8.3	8.0	7.7	7.7	7.9
ARMOR	8.0	7.0	7.3	8.0	7.6
PIRANHA (DC-1)	8.3	7.0	7.0	6.0	7.1
CHINOOK (H10G-OP)	8.0	7.7	7.3	5.3	7.1
007	7.3	7.3	7.0	6.0	6.9
V-8	7.7	7.3	7.3	5.3	6.9
CRYSTAL BLUE LINKS	6.7	7.7	7.0	5.7	6.8
PST-ORBS	7.3	7.7	7.0	5.0	6.8
SHARK	7.0	7.7	7.0	5.3	6.8
PC2.0 (PST-OCV6)	7.3	7.0	7.0	5.3	6.7
PENNCROSS	7.0	7.0	7.0	5.3	6.6
BARRACUDA	6.3	7.7	7.0	5.0	6.5
L-93XD	6.0	7.3	7.0	5.0	6.3
LSD VALUE	1.2	0.7	0.6	0.8	0.4
C.V. (%)	9.9	6.2	4.8	8.2	7.6

- 1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).
- 2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9C. GENETIC COLOR RATINGS OF COLONIAL BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME	IA1	IL1	MI1	WA1	MEAN
GREENTIME	7.0	6.7	6.7	6	6.6
MUSKET (PPG-AT 104)	7.0	6.0	6.3	6	6.3
DLFPS-AT/3026	6.3	6.0	6.0	6	6.1
LSD VALUE	1.4	0.5	0.8	0	0.4
C.V. (%)	13.0	5.4	7.4	0	8.3

- 1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).
- 2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10A. SPRING GREENUP RATINGS OF BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/				
NAME	IL1	MI1	NJ1	MEAN
CRYSTAL BLUE LINKS	7.3	6.7	7.0	7.0
CHINOOK (H10G-OP)	7.3	6.0	6.7	6.7
V-8	7.3	6.7	5.7	6.6
PIRANHA (DC-1)	7.7	3.7	8.0	6.4
GREENTIME	7.3	6.3	5.7	6.4
PC2.0 (PST-OCV6)	7.3	6.7	5.3	6.4
BARRACUDA	7.3	5.3	6.0	6.2
007	7.7	5.0	5.7	6.1
KINGDOM	8.0	6.3	4.0	6.1
ARMOR	8.0	6.0	4.0	6.0
MUSKET (PPG-AT 104)	7.3	6.0	4.7	6.0
L-93XD	7.0	4.0	6.7	5.9
PENNCROSS	7.0	6.7	4.0	5.9
SHARK	7.7	5.7	4.3	5.9
NIGHTLIFE	8.0	5.7	3.7	5.8
PST-ORBS	7.3	6.0	4.0	5.8
DLFPS-AT/3026	7.0	6.3	3.7	5.7
LSD VALUE	0.7	1.7	2.4	1.0
C.V. (%)	6.2	18.0	27.9	17.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10B. SPRING GREENUP RATINGS OF CREEPING BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME	IL1	MI1	NJ1	MEAN
CRYSTAL BLUE LINKS	7.3	6.7	7.0	7.0
CHINOOK (H10G-OP)	7.3	6.0	6.7	6.7
V-8	7.3	6.7	5.7	6.6
PIRANHA (DC-1)	7.7	3.7	8.0	6.4
PC2.0 (PST-OCV6)	7.3	6.7	5.3	6.4
BARRACUDA	7.3	5.3	6.0	6.2
007	7.7	5.0	5.7	6.1
KINGDOM	8.0	6.3	4.0	6.1
ARMOR	8.0	6.0	4.0	6.0
L-93XD	7.0	4.0	6.7	5.9
PENNCROSS	7.0	6.7	4.0	5.9
SHARK	7.7	5.7	4.3	5.9
NIGHTLIFE	8.0	5.7	3.7	5.8
PST-ORBS	7.3	6.0	4.0	5.8
LSD VALUE	0.7	1.7	2.3	1.0
C.V. (%)	6.2	18.0	26.6	16.9

- 1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).
- 2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10C. SPRING GREENUP RATINGS OF COLONIAL BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME	IL1	MI1	NJ1	MEAN
GREENTIME	7.3	6.3	5.7	6.4
MUSKET (PPG-AT 104)	7.3	6.0	4.7	6.0
DLFPS-AT/3026	7.0	6.3	3.7	5.7
LSD VALUE	0.8	1.8	2.6	1.1
C.V. (%)	6.5	17.8	35.0	19.4

- 1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).
- 2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11A. LEAF TEXTURE RATINGS OF BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME	NJ1
PIRANHA (DC-1)	8.0
MUSKET (PPG-AT 104)	7.3
BARRACUDA	7.0
DLFPS-AT/3026	7.0
007	6.7
L-93XD	6.7
CHINOOK (H10G-OP)	6.3
V-8	6.3
ARMOR	5.7
PST-ORBS	5.7
SHARK	5.7
CRYSTAL BLUE LINKS	5.0
GREENTIME	5.0
KINGDOM	5.0
NIGHTLIFE	5.0
PC2.0 (PST-OCV6)	4.7
PENNCROSS	3.7
LSD VALUE	1.1
C.V. (%)	11.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11B. LEAF TEXTURE RATINGS OF CREEPING BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME	NJ1
PIRANHA (DC-1)	8.0
BARRACUDA	7.0
007	6.7
L-93XD	6.7
CHINOOK (H10G-OP)	6.3
V-8	6.3
ARMOR	5.7
PST-ORBS	5.7
SHARK	5.7
CRYSTAL BLUE LINKS	5.0
KINGDOM	5.0
NIGHTLIFE	5.0
PC2.0 (PST-OCV6)	4.7
PENNCROSS	3.7
LSD VALUE	1.1
C.V. (%)	12.2

- 1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).
- 2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11C. LEAF TEXTURE RATINGS OF COLONIAL BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME	NJ1
MUSKET (PPG-AT 104)	7.3
DLFPS-AT/3026	7.0
GREENTIME	5.0
LSD VALUE	1.1
C.V. (%)	10.3

- 1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).
- 2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12A. FALL DENSITY RATINGS OF BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	NJ1
PIRANHA (DC-1)	7.3
CHINOOK (H10G-OP)	6.7
DLFPS-AT/3026	6.7
007	6.3
BARRACUDA	6.3
L-93XD	6.0
ARMOR	5.7
KINGDOM	5.7
MUSKET (PPG-AT 104)	5.7
V-8	5.7
SHARK	5.3
NIGHTLIFE	5.0
CRYSTAL BLUE LINKS	4.7
GREENTIME	4.7
PST-ORBS	4.7
PC2.0 (PST-OCV6)	4.0
PENNCROSS	4.0
LSD VALUE	1.3
C.V. (%)	14.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12B. FALL DENSITY RATINGS OF CREEPING BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	NJ1
PIRANHA (DC-1)	7.3
CHINOOK (H10G-OP)	6.7
007	6.3
BARRACUDA	6.3
L-93XD	6.0
ARMOR	5.7
KINGDOM	5.7
V-8	5.7
SHARK	5.3
NIGHTLIFE	5.0
CRYSTAL BLUE LINKS	4.7
PST-ORBS	4.7
PC2.0 (PST-OCV6)	4.0
PENNCROSS	4.0
LSD VALUE	1.2
C.V. (%)	13.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12C. FALL DENSITY RATINGS OF COLONIAL BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	NJ1
DLFPS-AT/3026	6.7
MUSKET (PPG-AT 104)	5.7
GREENTIME	4.7
LSD VALUE	1.9
C.V. (%)	20.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13A. DOLLAR SPOT RATINGS OF BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	IA1	MD1	NJ1	MEAN
CHINOOK (H10G-OP)	7.7	8.0	9.0	8.2
DLFPS-AT/3026	8.7	8.0	8.0	8.2
MUSKET (PPG-AT 104)	7.3	7.7	8.3	7.8
PIRANHA (DC-1)	8.0	6.3	9.0	7.8
L-93XD	8.3	7.3	7.0	7.6
007	8.0	6.7	7.3	7.3
BARRACUDA	8.3	5.3	6.7	6.8
V-8	7.0	6.0	7.3	6.8
CRYSTAL BLUE LINKS	7.7	6.7	5.7	6.7
KINGDOM	7.7	5.7	5.3	6.2
GREENTIME	7.0	6.0	4.7	5.9
NIGHTLIFE	7.3	4.7	4.7	5.6
SHARK	6.7	4.7	5.3	5.6
PENNCROSS	6.3	5.7	2.3	4.8
PC2.0 (PST-OCV6)	6.0	6.3	1.7	4.7
PST-ORBS	6.3	5.7	1.3	4.4
ARMOR	7.0	4.0	1.7	4.2
LSD VALUE	1.3	2.0	2.0	1.0
C.V. (%)	10.7	20.5	22.6	17.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13B. DOLLAR SPOT RATINGS OF CREEPING BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE 2/				
NAME	IA1	MD1	NJ1	MEAN
CHINOOK (H10G-OP)	7.7	8.0	9.0	8.2
PIRANHA (DC-1)	8.0	6.3	9.0	7.8
L-93XD	8.3	7.3	7.0	7.6
007	8.0	6.7	7.3	7.3
BARRACUDA	8.3	5.3	6.7	6.8
V-8	7.0	6.0	7.3	6.8
CRYSTAL BLUE LINKS	7.7	6.7	5.7	6.7
KINGDOM	7.7	5.7	5.3	6.2
NIGHTLIFE	7.3	4.7	4.7	5.6
SHARK	6.7	4.7	5.3	5.6
PENNCROSS	6.3	5.7	2.3	4.8
PC2.0 (PST-OCV6)	6.0	6.3	1.7	4.7
PST-ORBS	6.3	5.7	1.3	4.4
ARMOR	7.0	4.0	1.7	4.2
LSD VALUE	1.3	2.2	2.2	1.1
C.V. (%)	11.0	22.8	25.5	19.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13C. DOLLAR SPOT RATINGS OF COLONIAL BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE 2/				
NAME	IA1	MD1	NJ1	MEAN
DLFPS-AT/3026	8.7	8.0	8.0	8.2
MUSKET (PPG-AT 104)	7.3	7.7	8.3	7.8
GREENTIME	7.0	6.0	4.7	5.9
LSD VALUE	1.2	1.1	1.2	0.7
C.V. (%)	9.7	9.2	10.6	9.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14A. BROWN PATCH (WARM TEMPERATURE) RATINGS OF BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	IA1
ARMOR	9.0
CHINOOK (H10G-OP)	9.0
CRYSTAL BLUE LINKS	9.0
KINGDOM	9.0
L-93XD	9.0
NIGHTLIFE	9.0
PIRANHA (DC-1)	9.0
SHARK	9.0
V-8	9.0
BARRACUDA	8.7
PC2.0 (PST-OCV6)	8.7
DLFPS-AT/3026	8.3
PENNCROSS	8.3
007	8.0
GREENTIME	8.0
PST-ORBS	7.3
MUSKET (PPG-AT 104)	5.0
LSD VALUE	1.8
C.V. (%)	13.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14B. BROWN PATCH (WARM TEMPERATURE) RATINGS OF CREEPING BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	IA1
ARMOR	9.0
CHINOOK (H10G-OP)	9.0
CRYSTAL BLUE LINKS	9.0
KINGDOM	9.0
L-93XD	9.0
NIGHTLIFE	9.0
PIRANHA (DC-1)	9.0
SHARK	9.0
V-8	9.0
BARRACUDA	8.7
PC2.0 (PST-OCV6)	8.7
PENNCROSS	8.3
007	8.0
PST-ORBS	7.3
LSD VALUE	1.6
C.V. (%)	11.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14C. BROWN PATCH (WARM TEMPERATURE) RATINGS OF COLONIAL BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	IA1
DLFPS-AT/3026	8.3
GREENTIME	8.0
MUSKET (PPG-AT 104)	5.0
LSD VALUE	2.7
C.V. (%)	23.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15A. ANTHRACNOSE RATINGS OF BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

ANTHRACNOSE RATINGS 1-9; 9=NO DISEASE 2/

NAME	NJ1
DLFPS-AT/3026	9.0
GREENTIME	9.0
MUSKET (PPG-AT 104)	9.0
PC2.0 (PST-OCV6)	9.0
L-93XD	8.7
V-8	8.7
SHARK	8.0
ARMOR	7.3
CHINOOK (H10G-OP)	7.3
PIRANHA (DC-1)	7.0
BARRACUDA	6.7
NIGHTLIFE	6.7
007	6.3
KINGDOM	6.0
PST-ORBS	5.3
CRYSTAL BLUE LINKS	4.3
PENNCROSS	3.3
LSD VALUE	2.3
C.V. (%)	19.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15B. ANTHRACNOSE RATINGS OF CREEPING BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

ANTHRACNOSE RATINGS 1-9; 9=NO DISEASE 2/

NAME	NJ1
PC2.0 (PST-OCV6)	9.0
L-93XD	8.7
V-8	8.7
SHARK	8.0
ARMOR	7.3
CHINOOK (H10G-OP)	7.3
PIRANHA (DC-1)	7.0
BARRACUDA	6.7
NIGHTLIFE	6.7
007	6.3
KINGDOM	6.0
PST-ORBS	5.3
CRYSTAL BLUE LINKS	4.3
PENNCROSS	3.3
LSD VALUE	2.5
C.V. (%)	22.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15C. ANTHRACNOSE RATINGS OF COLONIAL BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

ANTHRACNOSE RATINGS 1-9; 9=NO DISEASE 2/

NAME	NJ1
DLFPS-AT/3026	9
GREENTIME	9
MUSKET (PPG-AT 104)	9
C.V. (%)	0
LSD VALUE	0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16A. FALL COLOR (NOVEMBER) RATINGS OF BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	IL1	NJ1	MEAN
CHINOOK (H10G-OP)	7.7	8.0	7.8
007	7.7	7.3	7.5
PIRANHA (DC-1)	7.0	8.0	7.5
L-93XD	7.3	6.7	7.0
BARRACUDA	7.7	6.0	6.8
V-8	7.0	6.3	6.7
KINGDOM	7.3	5.7	6.5
SHARK	7.7	5.3	6.5
ARMOR	7.0	5.7	6.3
NIGHTLIFE	7.7	5.0	6.3
PST-ORBS	7.7	4.7	6.2
CRYSTAL BLUE LINKS	7.3	5.0	6.2
PC2.0 (PST-OCV6)	7.3	4.7	6.0
PENNCROSS	8.0	3.7	5.8
DLFPS-AT/3026	7.7	3.7	5.7
GREENTIME	7.3	3.7	5.5
MUSKET (PPG-AT 104)	6.3	3.0	4.7
LSD VALUE	0.8	1.4	0.8
C.V. (%)	6.8	15.9	11.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16B. FALL COLOR (NOVEMBER) RATINGS OF CREEPING BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	IL1	NJ1	MEAN
CHINOOK (H10G-OP)	7.7	8.0	7.8
007	7.7	7.3	7.5
PIRANHA (DC-1)	7.0	8.0	7.5
L-93XD	7.3	6.7	7.0
BARRACUDA	7.7	6.0	6.8
V-8	7.0	6.3	6.7
KINGDOM	7.3	5.7	6.5
SHARK	7.7	5.3	6.5
ARMOR	7.0	5.7	6.3
NIGHTLIFE	7.7	5.0	6.3
PST-ORBS	7.7	4.7	6.2
CRYSTAL BLUE LINKS	7.3	5.0	6.2
PC2.0 (PST-OCV6)	7.3	4.7	6.0
PENNCROSS	8.0	3.7	5.8
LSD VALUE	0.8	1.5	0.8
C.V. (%)	6.5	15.8	11.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16C. FALL COLOR (NOVEMBER) RATINGS OF COLONIAL BENTGRASS CULTIVARS 1/
GROWN ON A FAIRWAY OR TEE
2017 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	IL1	NJ1	MEAN
DLFPS-AT/3026	7.7	3.7	5.7
GREENTIME	7.3	3.7	5.5
MUSKET (PPG-AT 104)	6.3	3.0	4.7
LSD VALUE	0.9	0.8	0.6
C.V. (%)	8.1	13.7	10.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

APPENDIX TABLE. SUMMARY OF TURFGRASS QUALITY RATINGS FOR BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE
2017 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF

NAME	QUALITY MEAN 1/	MAXIMUM IN TOP 25% 2/
007	6.5	11.1
ARMOR	6.0	22.2
BARRACUDA	6.5	44.4
CHINOOK (H10G-OP)	6.7	66.7
CRYSTAL BLUE LINKS	6.4	22.2
DLFPS-AT/3026	6.1	11.1
GREENTIME	5.6	11.1
KINGDOM	6.2	22.2
L-93XD	6.4	22.2
MUSKET (PPG-AT 104)	6.0	0.0
NIGHTLIFE	6.4	44.4
PC2.0 (PST-OCV6)	6.1	11.1
PENNCROSS	5.3	0.0
PIRANHA (DC-1)	6.7	22.2
PST-ORBS	5.9	11.1
SHARK	6.4	11.1
V-8	6.4	33.3
LSD VALUE	0.3	
C.V. (%)	8.0	

*/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD VALUE 0.05).

**/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

1/ MEAN AN AVERAGE OF ALL THE TURFGRASS QUALITY RATINGS FROM ALL LOCATIONS.

2/ MAXIMUM IN TOP 25%: THE PERCENTAGE OF LOCATIONS WHERE THAT ENTRY FINISHED IN THE TOP 25% OF ALL ENTRIES.